

Original Article

Secoemestrin C inhibits activation of NKT/conventional T cells and protects against concanavalin A-induced autoimmune hepatitis in mice

Xiaosheng Tan^{1,2*}, Lingjuan Sun^{1,2*}, Qin Li^{3*}, Changxing Qi³, Cheng Fu^{1,2}, Hucheng Zhu³, Xi Yang⁴, Hao Feng^{1,2}, Yakun Li^{1,2}, Yonghui Zhang³, Gang Chen^{1,2}

¹Institute of Organ Transplantation, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, Hubei Province, People's Republic of China; ²Key Laboratory of Organ Transplantation, Ministry of Education, NHC and Chinese Academy of Medical Sciences, People's Republic of China; ³Hubei Key Laboratory of Natural Medicinal Chemistry and Resource Evaluation, School of Pharmacy, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, People's Republic of China; ⁴Department of Pediatrics, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, People's Republic of China. *Equal contributors.

Received March 18, 2020; Accepted June 13, 2020; Epub July 15, 2020; Published July 30, 2020

Abstract: Secoemestrin C is an epitetrahydrodioxopiperazine isolated from *Aspergillus nidulans*, which has only been reported in terms of inhibiting tumor cell proliferation. Here, we determined the immunomodulatory and hepatoprotective effects of secoemestrin C in a mouse model of acute autoimmune hepatitis. In an *in vitro* assay, purified hepatic mononuclear cells (MNCs) from C57BL/6J mice were stimulated with concanavalin A (Con A) in the presence of secoemestrin C, and cell proliferation and cytokine production were measured. In an *in vivo* assay, mice with or without secoemestrin C pretreatment were injected with Con A to induce acute hepatitis. The secoemestrin C treatment dose-dependently suppressed cell proliferation and proinflammatory cytokine secretion in Con A-stimulated hepatic MNCs *in vitro*. In Con A-challenged mice, pre-injection with secoemestrin C significantly decreased the generation of proinflammatory cytokines and ameliorated liver injury. Furthermore, pretreatment with secoemestrin C significantly inhibited the Con A-induced activation of NKT and conventional T cells, and decreased IFN- γ production by these two cell populations. We conclude that secoemestrin C has an immunosuppressive effect on NKT and conventional T cells and has hepatoprotective activity in mouse autoimmune hepatitis. These findings provide new insights into the use of fungus-derived natural products for the treatment of autoimmune diseases.

Keywords: Secoemestrin C, concanavalin A, autoimmune hepatitis, NKT cells, conventional T cells, hepatoprotective effects

Introduction

Autoimmune hepatitis (AIH) is a chronic self-perpetuating liver inflammatory disease that affects all ages, both genders, and all ethnicities [1]. AIH may start with an episode of acute hepatitis and eventually lead to liver cirrhosis, liver cancer, the need for liver transplantation, or death [2]. The precise etiology of AIH is unknown, but it is been widely believed that natural killer T (NKT) cells and conventional T cells contribute to AIH-induced liver damage [3, 4]. The prominent features of these two cell populations during AIH include an accumulation in the liver, aberrant activation, and imbal-

anced Th1/Th2 responses [5, 6]. Once a T cell-mediated autoimmune response is initiated, hepatocytes can potentially be destroyed directly through Fas/Fas ligand (FasL) interactions and/or the action of TNF- α and perforins/granzyme B, or indirectly through the release of Th1-biased IFN- γ [3]. Therefore, T cells are expected to be the therapeutic targets in the treatment of AIH.

The mainstay of treatment for AIH is immunosuppression [7]. Concanavalin A (Con A)-induced acute hepatitis in mice is considered to be a well-characterized model for studying the pathogenesis of AIH and screening immunosup-




Figure 1. Secoemestrin C inhibits Con A-stimulated splenocyte proliferation and elevates SOCS proteins mRNA expression. (A) Chemical structure of secoemestrin C (SC) isolated from *Aspergillus nidulans*. (B, C) Splenocytes from C57BL/6J mice were stained with CFSE and stimulated with Con A ($2 \mu\text{g mL}^{-1}$) in the presence of secoemestrin C (SC) at the indicated concentration, and the cells were cultured for 72 h. Splenocyte proliferation was detected by flow cytometry. A representative histogram (B) and summary bar graph (C) are shown. (D) Splenocytes isolated from C57BL/6J mice were stimulated with Con A ($2 \mu\text{g mL}^{-1}$) in the presence of secoemestrin C (128 nM), and the cells were cultured for 24 h. The mRNA levels of SOCS1, SOCS2 and SOCS3 were measured by real-time PCR. Data are shown as means \pm SEM ($n=5$ per group). * $P<0.05$, ** $P<0.01$ and *** $P<0.001$ vs. the control group; ## $P<0.01$ and ### $P<0.001$ vs. the Con A-treated group.

pressive agents [8]. Con A primarily stimulates the recruitment and activation of NKT and conventional T cells, coinciding with the generation of large amounts of proinflammatory Th1 cytokines [9]. In recent years, based on the administration of Con A to model animals, many new therapeutic drugs, antibodies, and proteins have been developed to attenuate liver injury by inhibiting NKT/T cell activation and skewing [10].

Fungus-derived natural products and their derivatives, such as cyclosporine and mycophenolate mofetil, are widely used in the treatment of autoimmune diseases [11, 12]. Preparations of fungi and their metabolites have a low synthetic cost, high practical value, and broad application in medicine, thanks to their wide availability and high biological activity [13]. In a previous screening of anti-inflammatory agents, we found that secoemestrin C (Figure 1A), isolated from *Aspergillus nidulans*, had a potent inhibitory effect on Con A-induced splenocyte proliferation, suggesting potential immunosuppressive and anti-inflammatory activity. In the

present study, we sought to investigate whether secoemestrin C could protect against Con A-induced autoimmune hepatitis in mice and to explore potential mechanisms for such an effect.

Materials and methods


Experimental animals

Specific pathogen-free C57BL/6J (6-8 weeks, 18-23 g) female mice were purchased from Charles River Laboratories (Beijing, China) and maintained in controlled conditions (22°C, 50% humidity, 12 h light/dark cycle, with lights on at 7:00 AM). The mouse care and experimental protocols conducted in this study were approved by the Huazhong University of Science and Technology Animal Care and Use Committee.

Isolation and identification of secoemestrin C

The fungus *Aspergillus nidulans* was growing on rice culture medium (40 kg) at room temperature for 21 days. The fermented medium

Table 1. ^1H (400 MHz) and ^{13}C (100 MHz) nuclear magnetic resonance (NMR) data of secoemestrin C (CDCl_3 , δ in ppm, J in Hz)



NO.	δ_{H} (J in Hz)	δ_{C}	DEPT
1		167.2	C
3	5.20 s	67.6	CH
4		163.8	C
5a	5.46 dd (8.4, 2.4)	60.0	CH
6	5.31 dt (9.2, 2.7)	72.0	CH
7	4.87 dd (8.4, 1.9)	107.6	CH
8	6.30 dd (8.4, 2.4)	139.3	CH
10	6.85 d (2.4)	142.9	CH
10a		110.4	C
11	4.89 s	78.1	CH
11a		80.0	C
1'		123.1	C
2'	8.09 d (2.0)	124.0	CH
3'		144.3	C
4'		155.3	C
5'	7.07 d (8.6)	112.1	CH
6'	8.06 dd (8.6, 2.0)	129.3	CH
7'		165.3	C
1''		145.8	C
2''		153.4	C
3''	7.13 d (8.3)	116.3	CH
4''	7.54 dd (8.3, 2.0)	129.1	CH
5''		129.9	C
6''	7.56 d (2.0)	118.1	CH
7''	9.77 s	190.9	CH
2-NMe	3.02 s	32.6	CH_3
4'-OMe	3.92 s	56.3	CH_3

was then extracted with 95% ethanol, and the extract was suspended in water and further treated with ethyl acetate (AcOEt) to obtain a dark brown extract (300 g). The AcOEt extract was chromatographed using silica gel (100-200 mesh) and eluted with petroleum ether/

AcOEt (50:1-0:100) to give six fractions (Fr1-Fr6). Secoemestrin C (5.2 g) was isolated from Fraction 4, and the purity ($\geq 99.0\%$) was determined via HPLC with a UV detector. The structure of secoemestrin C was elucidated based on NMR and HRESIMS (Table 1; Figures 1A and 2) and found to be identical to that reported previously [14].

Reagents and antibodies

Carboxy-fluorescein diacetate succinimidyl ester (CFSE) was purchased from Invitrogen (ThermoFisher Scientific). Concanavalin A type IV from *Canavalia ensiformis* (C2010) was obtained from Sigma-Aldrich. Alanine aminotransferase (ALT), aspartate aminotransferase (AST), and lactate dehydrogenase (LDH) were measured with an automated biochemical analyzer BS-200 (Mindray, China). GolgiStop brefeldin A solution and Fix/Perm Buffer Set were obtained from Biolegend. Flow cytometry antibodies: antibodies against mouse TCR β (H57-597), NK1.1 (PK136), CD4 (GK1.5), CD8 (53-6.7), CD69 (H1.2F3), and IFN- γ (XMG1.2) were purchased from BD Biosciences and Biolegend, with fluorescence conjugation. Cytokines in serum or in culture supernatants were determined by using a cytometric bead array-based mouse Th1/Th2/Th17 Cytokine Kit (BD Biosciences).

Isolation of hepatic mononuclear cells and splenocytes

Hepatic mononuclear cells (MNCs) and splenocytes were isolated as previously described [15]. In brief, each mouse liver was perfused with phosphate buffer saline to eliminate blood, and the liver tissue was removed and pressed through a 70- μm cell strainer (BD Biosciences). The liver cell suspension was then collected. Hepatic MNCs were purified by density gradient centrifugation in a 38% Percoll solution (GE Healthcare). Purified splenocytes were prepared by homogenizing with a syringe, followed by passage through a 0.1-mm sterile nylon mesh and erythrocyte depletion.

Real-time PCR

To determine mRNA levels of the suppressors of cytokine signaling (SOCS) family of proteins, total RNA was extracted from liver tissues using RNAfast200 (Fastagen, Shanghai,




Figure 2. High resolution electrospray ionization mass spectroscopy (HRESIMS) spectrum of secoemestrin C. The compound was dissolved in methanol. Data were acquired by using a Bruker microTOF II spectrometer.

China) according to the manufacturer's instructions; RNA was reverse-transcribed with a PrimeScript RT Reagent Kit (Takara) to generate cDNA. The PCR reaction mixture was prepared using a SYBR Green Premix Ex Taq II Kit (Takara) with the primers as follows: SOCS1 (forward: 5'-CTGCGGCTTCTATTGGGGAC-3'; reverse: 5'-AAAAGGCAGTCGAAGGTCTCG-3'); SOCS2 (forward: 5'-AGTTCGCATTCAGACTACCTACT-3'; reverse: 5'-TGGTACTCAATCCGCAGGTTAG-3'); SOCS3 (forward: 5'-ATGGTCACCCACAGCAAGTTT-3'; reverse: 5'-TCCAGTAGAATCCGCTCTCCT-3'); and HPRT1 (forward: 5'-AGTACAGCCCCAAAATGGTTA-3'; reverse: 5'-CTTAGGCTTTGTATTGGCTT-3').

Hepatitis induction and treatment

C67BL/6J mice in the Con A-challenged and secoemestrin C-treated groups were injected intravenously with a single dose of Con A (12 mg·kg⁻¹) to simulate autoimmune hepatitis. In the case of the secoemestrin C-treated group, secoemestrin C (1 mg·kg⁻¹ or 10 mg·kg⁻¹) was intraperitoneally injected for 3 days, and then the mice were injected with Con A 1 h after the last secoemestrin C injection.

Blood samples were collected from the orbital venous plexus under mild pentobarbital anesthesia 8 h after the Con A injection, and the serum was separated by centrifugation. Later, mice were sacrificed by cervical disloca-

tion under mild pentobarbital anesthesia, and livers were harvested.

Cytokine assay

The levels of IL-2, IL-4, IL-6, IL-10, IL-17, IFN- γ , and TNF- α in the serum 8 h after Con A challenge or in culture supernatants 72 h after Con A stimulation were measured with a CBA kit for mouse Th1/Th2/Th17 cytokines.

Flow cytometry

Procedures for surface staining, intracellular cytokine staining, and CBA kit use were performed according to the manufacturer's instructions.

Flow cytometry data were acquired on a BD FACSCelesta and analyzed with FlowJo software (TreeStar).

Histological examination

Liver samples were fixed in 4% formaldehyde and dehydrated with xylene, absolute ethyl alcohol, and 75% alcohol, then embedded in paraffin wax. Four micron-thick sections were cut and stained with hematoxylin-eosin (H&E) and observed by three pathologists in a blinded manner. The extent of the pathology was scored from 0 (no pathology) to 3 (severe pathology) as previously described [16].

Immunohistochemistry and immunofluorescent staining

Liver sections were stained with primary antibody (anti-Fas or anti-cleaved caspase-3) according to a standard protocol (Servicebio, China). Apoptotic cells were evaluated using a TUNEL assay kit (Roche Applied Science) according to the manufacturer's instructions.

Statistical analysis

Graphs were generated and analyzed by GraphPad Prism 6.0 (GraphPad Software). Data were analyzed by Student's *t*-test for comparisons of groups with normal distribution and by equal variance or Mann-Whitney *U* test for non-normally distributed variables among groups.




Figure 3. Inhibitory effect of secoemestrin C on the proliferation and cytokine production of Con A-stimulated liver mononuclear cells (MNCs). Liver MNCs from C57BL/6J mice were stimulated with $2 \mu\text{g/mL}$ Con A in the presence of secoemestrin C (SC) at the indicated concentrations. After 72 h of culture, hepatic MNC proliferation was determined by CFSE-dilution assay. Cytokine concentrations in culture supernatants were quantified by cytometric bead array kits. (A) The gating strat-

Secoemestrin C ameliorates Con A-induced hepatitis

egy for hepatic NKT and conventional T cells is displayed. (B) Bar graphs depict the percentages of TCR β ⁺NK1.1⁺ NKT cells (NKT) and TCR β ⁺NK1.1⁻ conventional T cells (conT) in liver MNCs. (C, D) A representative histogram (C) and summary bar graph (D) depict the percentages of proliferated cells. (E-J) Levels of IFN- γ (E), IL-2 (F), IL-6 (G), IL-17 (H), TNF- α (I), and IL-4 (J) for the various groups are shown in the bar graphs. Bar graphs indicate means \pm SEM (n=5). **P<0.01 and ***P<0.001 vs. the control (NC) group. #P<0.05 and ##P<0.01 vs. the Con A-treated group.

Data are presented as means \pm standard error of the mean (SEM). P<0.05 was considered significant.

Results

Secoemestrin C suppresses the proliferation of Con A-stimulated splenocytes

A total of 5.2 g of secoemestrin C was isolated from *Aspergillus nidulans*, which had been growing on rice culture medium (40 kg). The structure of the compound was elucidated by NMR and HR-ESI-MS (Figures 1A and 2; Table 1). The purity of the secoemestrin C was \geq 99.0%, as determined via high performance liquid chromatography with a UV detector. To investigate whether secoemestrin C has the potential to suppress immunological reactions, a Con A-induced splenocyte proliferation assay was carried out in the presence of secoemestrin C. Mouse splenocytes strongly proliferated in response to the Con A stimulation *in vitro* (Figure 1B, 1C). However, secoemestrin C treatment potently inhibited this Con A-stimulated splenocyte proliferation (Figure 1B, 1C). Considering the important role of SOCS proteins in the regulation of autoimmunity [17, 18], we further detected the mRNA expression of SOCS1, SOCS2 and SOCS3 in splenocytes. Secoemestrin C treatment directly induced significant up-regulated expression of these SOCS proteins in splenocytes. Compared with untreated control splenocytes, the mRNA expression of SOCS1, SOCS2 and SOCS3 in Con A-stimulated splenocytes was significantly decreased, while secoemestrin C treatment could significantly up-regulate the expression of these SOCS proteins in Con A-stimulated splenocytes (Figure 1D).

Secoemestrin C inhibits the proliferation of liver mononuclear cells (MNCs) and the production of cytokines induced by Con A in vitro

Tissue-resident immune cells play a key role in local and systemic immune responses. To determine whether secoemestrin C can also inhibit the activation of liver-resident lympho-

cytes, liver MNCs were isolated from normal mice and stimulated by Con A *in vitro*. FACS analysis showed that the isolated hepatic MNCs included 30.74 \pm 4.08% natural killer T (NKT) cells and 23.78 \pm 1.18% conventional T cells (Figure 3A, 3B). In response to Con A stimulation, hepatic MNCs strongly proliferated and released large amounts of cytokines, including IFN- γ , IL-2, IL-6, IL-17, TNF- α , and IL-4 (Figure 3C-J). Interestingly, secoemestrin C treatment significantly attenuated Con A-stimulated liver MNC proliferation and cytokine production in a dose-dependent manner (Figure 3C-H, 3J). Above all, secoemestrin C was able to restrain hepatic lymphocyte proliferation and cytokine production *in vitro*, which may enable it to exert an *in vivo* protective effect on Con A-induced and T cell-mediated autoimmune hepatitis in mice.

Secoemestrin C ameliorates liver injury in Con A-challenged mice

Next, we evaluated the effects of secoemestrin C on Con A-induced hepatitis *in vivo*. A severe liver injury was observed in mice 8 h after Con A administration, which manifested itself as a significant elevation in serum levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), and lactate dehydrogenase (LDH) when compared to the normal (control) mice (Figure 4A-C). In contrast, pretreatment with secoemestrin C at a higher dose (10 mg·kg⁻¹) almost completely inhibited the increase in the serum levels of ALT, AST, and LDH induced by the Con A challenge (Figure 4A-C).

To further confirm the hepatoprotective effect of secoemestrin C, hematoxylin-eosin (H&E)-stained liver tissue sections were evaluated for histopathology. In the Con A-treated group, massive bridging coagulative necrosis of hepatocytes was visualized by light microscopy (Figure 4D). However, the hepatic pathological changes induced by Con A were dramatically reduced in the mice treated with high-dose secoemestrin C (Figure 4D, 4E).

Secoemestrin C ameliorates Con A-induced hepatitis




Figure 4. Protective effect of secoemestrin C in Con A-induced liver injury. Secoemestrin C (SC) was dissolved in DMSO and further diluted with normal saline. Female C57BL/6J mice were pretreated by intraperitoneal injection with SC at 10 mg·kg⁻¹ (SC₁₀) or 1 mg·kg⁻¹ on days -2, -1, and 1 h before i.v. injection with Con A (12 mg/kg). Mice without Con A injection served as controls (NC). (A-C) Serum levels of alanine aminotransferase (ALT), aspartate aminotransferase (AST), and lactate dehydrogenase (LDH) were measured 8 h after Con A injection. Data shown represent means ± SEM (n=6 per group). (D) Liver sections were stained with hematoxylin and eosin (H&E). The arrow indicates hepatic ecchymosis; the star indicates infiltrated leukocytes, and the triangle shows an area of necrosis (magnification: ×200). (E) Histopathologic scores were evaluated, and the results are presented as the means ± SEM for three mice per group. (F and G) Liver sections were stained with TUNEL staining. Representative images (F) and a bar graph of apoptotic area (G) are shown (magnification: ×200). Data represent the mean ± SEM. **P<0.01 and ***P<0.001 vs. the control (NC) group; ###P<0.01 and ###P<0.001 vs. the Con A-treated group.

The apoptotic response plays an important role in Con A-induced liver injury. To investigate the pathway of secoemestrin C involved in the regulation of hepatocyte apoptosis, we used TUNEL staining as well as Fas and cleaved caspase-3 staining to analyze hepatocyte apoptosis in the livers of the mice. The area of apoptotic cells in the Con A group was significantly larger than that in the normal control group (Figure 4F, 4G). Mice pretreated with a high dose of secoemestrin C showed a much smaller apoptotic area in the liver (Figure 4F, 4G). Furthermore, the expression of two apoptosis-related proteins, Fas and cleaved caspase-3, was markedly elevated in Con A-challenged mice, whereas expression of the two proteins

was almost completely suppressed in the mice who received the high-dose secoemestrin C treatment (Figure 5). Together, these results indicate that pretreatment with secoemestrin C provides significant protection against Con A-induced hepatic damage.

Secoemestrin C reduces serum levels of proinflammatory cytokines in mice with Con A-induced hepatitis

A balance between proinflammatory and anti-inflammatory factors is essential for maintaining liver homeostasis. Proinflammatory cytokines play a critical role in the initiation and propagation of the autoimmune response. To investigate the effect of secoemestrin C treat-




Figure 5. Secoemestrin C reduces hepatic Fas and cleaved caspase-3 expression in Con A induced liver injury. Secoemestrin C (SC) was dissolved in DMSO and further diluted with normal saline. Female C57BL/6J mice were intraperitoneally pretreated with SC 10 mg·kg⁻¹ (SC₁₀) or vehicle (normal control, NC) on days -2, -1, and 1 h before i.v. injection with Con A (12 mg·kg⁻¹). (A and B) Liver tissue sections were stained with Fas. Representative images (A) and bar graph (B) of Fas positive area percentage are shown (magnification: ×200). (C and D) Cleaved caspase-3 was stained and the ratio of apoptotic nuclei to total nuclei (the apoptosis rate) was calculated. Representative images (C) and bar graph (D) of cleaved caspase 3 positive cells per 103 cells for each group are shown (magnification: ×400). Data are presented with mean ± SEM (three mice per group). **P<0.01 and ***P<0.001 vs. control group; ##P<0.01 and ###P<0.001 vs. Con A group.

ment on the generation of inflammatory cytokines in mice with Con A-induced liver injury, we examined the serum levels of IFN- γ , TNF- α , IL-2, IL-4, IL-6, and IL-17 using a cytometric bead array kit. In the Con A-challenged group, we found that the levels of IFN- γ , IL-2, TNF- α , IL-17, and IL-6 were dramatically increased when compared to those in the normal control mice. However, pretreatment with secoemestrin C significantly suppressed the production of these cytokines induced by Con A (Figure 6A-E). There were no significant differences in the serum level of IL-4 among the groups (Figure 6F).

Secoemestrin C inhibits Con A-induced activation of NKT cells and conventional T cells in liver

The recruitment and activation of NKT cells and conventional T cells contribute remarkably to the development and progression of Con A-induced hepatic inflammation. To investigate whether secoemestrin C has any effect on lymphocyte subsets, we analyzed liver MNCs 8 h after Con A injection. As compared to the normal control group, the percentage of hepatic NKT cells in Con A-challenged mice was significantly decreased, whereas the percentages of




Figure 6. Secoemestrin C treatment inhibits the production of serum proinflammatory cytokines in Con A-challenged mice. Serum was prepared from blood collected 8 h after Con A injection. Serum levels of IFN-γ (A), IL-2 (B), IL-6 (C), IL-17 (D), TNF-α (E), and IL-4 (F) were quantified by cytometric bead array kit. Bar graphs depict the concentrations of the indicated cytokines from the various groups. Values are presented as means \pm SEM (six mice per group). **P<0.01 vs. the control (NC) group. #P<0.05 and ##P<0.01 vs. the Con A-treated group.




Figure 7. Secoemestrin C rescues hepatic NKT decrease in Con A-challenged mice. C57BL/6J mice were sacrificed 8 h after being injected with Con A, and liver mononuclear cells (MNCs) were collected. The cells were stained with mAbs against TCRβ, CD4, CD8, or NK1.1, then analyzed by flow cytometry.

etry. A. Bar graphs depict the percentages of TCRβ⁺NK1.1⁺ NKT cells (NKT) vs. MNCs and of TCRβ⁺NK1.1⁻ conventional T cells (conT) vs. MNCs. B. Summary bar graphs display conventional CD4⁺ and CD8⁺ T cell (CD4⁺ conT and CD8⁺ conT) proportions in conventional T cells. Values shown are means \pm SEM (five mice per group). ***P<0.001 vs normal the control (NC) group. ##P<0.01 vs. the Con A-treated group.

conventional CD4⁺ and CD8⁺ T cells in the liver were relatively stable (Figure 7). Notably, secoemestrin C treatment largely rescued hepatic NKT cell depletion, as evidenced by a significantly increased ratio of hepatic NKT cells to liver MNCs in the treatment group when compared to that of the Con A-challenged group (Figure 7A).

To determine whether secoemestrin C can suppress the




Figure 8. Inhibitory effects of secoemestrin C on hepatic NKT cell and conventional T cell activation and proinflammatory cytokine secretion in Con A-challenged mice. Mice were sacrificed 8 h after Con A injection, and hepatic MNCs were prepared. (A, B) NKT cells as well as CD4⁺ and CD8⁺ conventional T cells were analyzed by flow cytometry for CD69 expression. Histogram with numbers (A) shows the mean fluorescence intensity (MFI) of CD69 on every subset of the normal control (NC), Con A-treated, and Con A+SC₁₀-treated groups. A summary bar graph (B) displays the CD69 levels on NKT cells as well as CD4⁺ and CD8⁺ conventional T cells. (C, D) Liver MNCs were cultured in medium with GolgiStop for a further 4 h, and intracellular staining for IFN-γ was performed. Representative dot plots (C) show IFN-γ expression by NKT cells (left panel), CD4⁺ conventional T cells (CD4⁺ conT, middle panel), and CD8⁺ conventional T cells (CD8⁺ conT, right panel). Bar graphs (D) depict the percentages of IFN-γ-producing cells in the corresponding liver MNC subsets. Data are shown as means ± SEM (five mice per group). *P<0.05, **P<0.01, and ***P<0.001 vs. the normal control (NC) group. #P<0.05 and ##P<0.01 vs. the Con A-treated group.

activation of hepatic NKT cells and conventional T cells *in vivo*, we measured the expression of CD69 and IFN-γ by FACS analysis. Secoemestrin C treatment significantly inhibited the increase in CD69 expression on hepatic NKT cells and conventional CD4⁺ and CD8⁺ T

cells that was seen in the Con A-challenged group (Figure 8A, 8C). In addition, treatment with secoemestrin C also down-regulated the elevated expression of IFN-γ in hepatic NKT and conventional CD4⁺ T cells after Con A challenge *in vivo* (Figure 8B, 8D).

Discussion

Fungi are widely distributed across the globe and have been identified as a rich source of new pharmacologically active ingredients. The members of the *Aspergillus* genus are already an important source of novel pharmacological metabolites such as alkaloids, meroterpenoids, and lignans. Secoemestrin C is an epitetrahydrodioxopiperazine extracted from *Aspergillus nidulans*. The biological activity of secoemestrin C and its analog has only been reported in terms of inhibiting tumor cell proliferation [19]. Here, we first present evidence to show that secoemestrin C can inhibit Con A-induced mouse splenocyte proliferation as well as hepatic MNC activation and proliferation *in vitro*, and then we demonstrate that this molecule has a protective effect *in vivo* on Con A-induced autoimmune hepatitis in mice.

Accumulating evidence suggests that CD4⁺ T cells [10, 20], CD8⁺ T cells [21], and NKT cells [22, 23] are critical components of Con A-induced liver damage. After Con A challenge, NKT cells and conventional T cells are activated and either directly participate in perforin/granzyme- and/or Fas/Fas ligand-mediated cytotoxicity or indirectly release large amounts of proinflammatory cytokines, thus leading to hepatocyte death. Therefore, agents that can inhibit the activation of T cells and/or NKT cells are most likely capable of alleviating hepatitis induced by Con A. In our *in vitro* experiments, we found that secoemestrin C can potently suppress the proliferation of splenocytes and activation of hepatic MNCs induced by Con A, indicating that it may have a significant protective effect on Con A-induced liver damage. As predicted, pretreatment with secoemestrin C markedly protected against liver injury in Con A-challenged mice, as evidenced by decreased serum levels of liver enzymes and proinflammatory cytokines, attenuated bridging coagulative necrosis of hepatocytes, and a lessened apoptotic response in the livers. In addition, the levels of serum transaminase and hepatocyte apoptosis in secoemestrin C-treated mice were similar to those in normal control mice, indicating that secoemestrin C has limited hepatic toxicity. Thus, secoemestrin C appears to be a good prospect for clinical application in immunosuppression.

In addition to aggravating inflammation and liver injury, activation of NKT cells also leads to Fas/FasL-mediated apoptotic elimination of NKT cells during Con A challenge [24]. Our results have confirmed the rapid disappearance of activated NKT cells in Con A-challenged livers. Although we did not detect any expression of FasL on NKT cells, our observation of a lowered CD69 expression level, together with a restored NKT cell ratio, suggest that the activation-induced cell death (AICD) of NKT cells is largely abolished in the secoemestrin C-treated mice.

Given that the depletion of NKT cells begins as early as 4 h after Con A administration [24], it appears that an NKT cell-derived proinflammatory cytokine response, rather than NKT cell-mediated cytotoxicity, plays a particularly important role in the latter phase of inflammation induced by Con A. In the treatment of autoimmune disease, targeting the production of proinflammatory cytokines has been approved as a potential therapy [25, 26]. It has been reported that SOCS3 suppresses IFN- γ production from NKT and T cells, thereby protecting against Con A-induced liver injury [27]. Our results show that secoemestrin C can down-regulate IFN- γ production by NKT and conventional T cells after Con A challenge. In addition, we have found that secoemestrin C is able to suppress cytokine production by Con A-stimulated hepatic MNCs. Suppressors of cytokine signaling (SOCS) proteins, especially SOCS1, SOCS2 and SOCS3 have been reported to play essential roles in inflammation and autoimmunity [17, 18]. In the present study, pretreatment of secoemestrin C was found to significantly elevate the mRNA expression of SOCS1, SOCS2 and SOCS3 in splenocytes with or without Con A stimulation, suggesting that secoemestrin C may have the capability to negatively regulate the cytokine-JAK-STAT pathway by facilitating the production of SOCS proteins. The regulatory mechanism(s) by which secoemestrin C inhibits proinflammatory cytokine production remain to be further explored.

Overall, we have demonstrated for the first time that secoemestrin C can inhibit NKT- and conventional T-cell activation stimulated by Con A and thus can protect against Con A-induced autoimmune hepatitis in mice. Our results provide evidence for the immunosup-

pressive activity of secoemestrin C and offer novel insights regarding the development of fungi-derived metabolites for the treatment of T cell-dependent autoimmune diseases.

Acknowledgements

This work was supported by National Nature Science Foundation of China (No. 81801587); project funded by China Postdoctoral Science Foundation (No. 2019M662642). We thank Dr. Deborah McClellan for editorial assistance.

Disclosure of conflict of interest

None.

Address correspondence to: Dr. Gang Chen, Institute of Organ Transplantation, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, 1095 Jiefang Avenue, Wuhan 430030, People's Republic of China. Tel: +86-27-83662892; Fax: +86-27-83662892; E-mail: gchen@tjh.tjmu.edu.cn; Dr. Yonghui Zhang, Hubei Key Laboratory of Natural Medicinal Chemistry and Resource Evaluation, Tongji Medical College, Huazhong University of Science and Technology, 13 Hangkong Road, Wuhan 430030, People's Republic of China. Tel: +86-27-83692892; Fax: +86-27-83692892; E-mail: zhangyh@mails.tjmu.edu.cn

References

- [1] Mieli-Vergani G, Vergani D, Czaja AJ, Manns MP, Krawitt EL, Vierling JM, Lohse AW and Montano-Loza AJ. Autoimmune hepatitis. *Nat Rev Dis Primers* 2018; 4: 18017.
- [2] Manns MP, Lohse AW and Vergani D. Autoimmune hepatitis - update 2015. *J Hepatol* 2015; 62 Suppl: S100-11.
- [3] Mieli-Vergani G, Vergani D, Czaja AJ, Manns MP, Krawitt EL, Vierling JM, Lohse AW and Montano-Loza AJ. Autoimmune hepatitis. *Nat Rev Dis Primers* 2018; 4: 18017.
- [4] Mattner J. Natural killer T (NKT) cells in autoimmune hepatitis. *Curr Opin Immunol* 2013; 25: 697-703.
- [5] Senaldi G, Portmann B, Mowat AP, Mieli-Vergani G and Vergani D. Immunohistochemical features of the portal tract mononuclear cell infiltrate in chronic aggressive hepatitis. *Arch Dis Child* 1992; 67: 1447-1453.
- [6] Santodomingo-Garzon T and Swain MG. Role of NKT cells in autoimmune liver disease. *Autoimmun Rev* 2011; 10: 793-800.
- [7] Vergani D and Mieli-Vergani G. Cutting Edge issues in autoimmune hepatitis. *Clin Rev Allergy Immunol* 2012; 42: 309-321.
- [8] Tiegs G, Hentschel J and Wendel A. A T cell-dependent experimental liver injury in mice inducible by concanavalin A. *J Clin Invest* 1992; 90: 196-203.
- [9] Heymann F, Hamesch K, Weiskirchen R and Tacke F. The concanavalin A model of acute hepatitis in mice. *Lab Anim* 2015; 49: 12-20.
- [10] Wang HX, Liu M, Weng SY, Li JJ, Xie C, He HL, Guan W, Yuan YS and Gao J. Immune mechanisms of concanavalin A model of autoimmune hepatitis. *World J Gastroenterol* 2012; 18: 119-125.
- [11] Lee CL, Sit WH, Jiang PP, So IW and Wan JM. Polysaccharopeptide mimics ciclosporin-mediated Th1/Th2 cytokine balance for suppression of activated human T cell proliferation by MAPKp38 and STAT5 pathways. *Pharm Pharmacol* 2008; 60: 1491-1499.
- [12] Fakih R, Matiello M, Chitnis T and Stankiewicz JM. Efficacy and safety of mycophenolate mofetil in progressive multiple sclerosis patients. *J Neurol* 2018; 265: 2688-2694.
- [13] Sharma M and Sharma R. Drugs and drug intermediates from fungi: striving for greener processes. *Crit Rev Microbiol* 2016; 42: 322-338.
- [14] Ooiike M, Nozawa K and Kawai KI. An epithiodioxopiperazine related to emestrin from *Emericella foveolata*. *Phytochemistry* 1997; 46: 123-126.
- [15] Zhou YQ, Weng XF, Dou R, Tan XS, Zhang TT, Fang JB and Wu XW. Betulin from *Hedyotis hedyotideae* ameliorates concanavalin A-induced and T cell-mediated autoimmune hepatitis in mice. *Acta Pharmacol Sin* 2017; 38: 201-210.
- [16] Yan JB, Jie ZL, Hou LF, Wanderley JL, Soong L, Gupta S, Qiu SM, Chan TS and Sun JR. Parenchymal expression of CD40 exacerbates adenovirus-induced hepatitis in mice. *Hepatology* 2011; 53: 1455-1467.
- [17] Liang Y, Xu WD, Peng H, Pan HF and Ye DQ. SOCS signaling in autoimmune diseases: molecular mechanisms and therapeutic implications. *Eur J Immunol* 2014; 44: 1265-1275.
- [18] Yoshimura A, Suzuki M, Sakaguchi R, Hanada T and Yasukawa H. SOCS, inflammation, and autoimmunity. *Front Immunol* 2012; 3: 20.
- [19] Onodera H, Hasegawa A, Tsumagari N, Nakai R, Ogawa T and Kanda Y. MPC1001 and its analogues: new antitumor agents from the fungus *Cladophium* species. *Org Lett* 2004; 6: 4101-4104.
- [20] Mizuhara H, O'Neil E, Seki N, Ogawa T, Kusunoki C, Otsuka K, Satoh S, Niwa M, Senoh H and Fujiwara H. T cell activation-associated hepatic injury: mediation by tumor necrosis factors and protection by interleukin 6. *J Exp Med* 1994; 179: 1529-1537.

Secoemestrin C ameliorates Con A-induced hepatitis

- [21] Watanabe Y, Morita M and Akaike T. Concavalin A induces perforin-mediated but not Fas-mediated hepatic injury. *Hepatology* 1996; 24: 702-710.
- [22] Kumar V. NKT-cell subsets: promoters and protectors in inflammatory liver disease. *J Hepatol* 2013; 59: 618-620.
- [23] Takeda K, Hayakawa Y, Van Kaer L, Matsuda H, Yagita H and Okumura K. Critical contribution of liver natural killer T cells to a murine model of hepatitis. *Proc Natl Acad Sci U S A* 2000; 97: 5498-5503.
- [24] Takeda K, Hayakawa Y, Van Kaer L, Matsuda H, Yagita H and Okumura K. Critical contribution of liver natural killer T cells to a murine model of hepatitis. *Proc Natl Acad Sci U S A* 2000; 97: 5498-5503.
- [25] Salvi V, Gianello V, Tiberio L, Sozzani S and Bosio D. Cytokine targeting by miRNAs in autoimmune diseases. *Front Immunol* 2019; 10: 15.
- [26] Danese S, Argollo M, Le Berre C and Peyrin-Biroulet L. JAK selectivity for inflammatory bowel disease treatment: does it clinically matter? *Gut* 2019; 68: 1893-1899.
- [27] Nakaya M, Hashimoto M, Nakagawa R, Wabayashi Y, Ishizaki T, Takada I, Komai K, Yoshida H and Yoshimura A. SOCS3 in T and NKT cells negatively regulates cytokine production and ameliorates Con A-induced hepatitis. *J Immunol* 2009; 183: 7047-7053.